

Table 3G
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Metro Area

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dane Cty	2,147	41.5	20.9	26.0	0.4	8.7	2.6	0.0
Mil. Cty	10,433	18.2	50.2	25.6	0.5	3.9	1.5	0.0
Racine	1,475	37.3	28.9	31.1	0.5	0.6	1.6	0.0
Rest of WI	20,879	73.6	3.0	14.2	3.0	4.6	1.7	0.0
Wisconsin	34,934	53.6	19.3	19.1	2.0	4.5	1.7	0.0
Nation (Prior Year)	1,153,475	41.5	22.7	28.7	1.1	3.1	0.9	2.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4G
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Metro Area

Page: 2
Run Date: 1 / 10 / 09

Comparison of Age Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dane Cty	2,147	0.4	5.8	12.1	62.2	18.5	1.0	0.0
Mil. Cty	10,433	0.6	8.6	14.3	58.6	16.5	1.3	0.0
Racine	1,475	0.3	8.2	15.5	60.5	14.6	0.9	0.0
Rest of WI	20,879	0.2	5.5	13.7	63.5	16.0	1.1	0.1
Wisconsin	34,934	0.3	6.6	13.8	61.8	16.2	1.2	0.1
Nation (Prior Year)	1,153,475	0.4	6.5	12.6	60.0	19.0	1.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5G
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by Metro Area

Comparison of Maternal Health Indicators (2, 3)																	
Metro Area	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum								
	Number	% (Rank)	%		Number	% (Rank)	% (Rank)	Number	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Dane Cty	2,122	10.5 (4)	42.6 (1)		1,289	27.3 (4)	43.8 (2)	404	27.0 (1)		1,441	21.0 (1)		1,902	4.4 (3)	1,902	5.1 (2)
Mil. Cty	10,333	8.3 (1)	49.9 (3)		7,101	26.7 (3)	43.3 (1)	1,770	43.6 (4)		8,921	34.8 (4)		9,361	4.3 (2)	9,361	7.1 (4)
Racine	1,459	9.4 (2)	51.9 (4)		1,279	25.6 (2)	45.6 (3)	228	32.9 (3)		1,265	25.0 (3)		1,254	4.9 (4)	1,254	6.9 (3)
Rest of WI	20,590	10.4 (3)	44.6 (2)		14,495	23.4 (1)	46.1 (4)	3,408	28.5 (2)		17,511	21.6 (2)		18,644	3.9 (1)	18,644	4.8 (1)
Wisconsin	34,504	9.7	46.4		24,164	24.7	45.1	5,810	33.1		29,138	25.8		31,161	4.1	31,161	5.6
Nation (Prior Year)	1,081,550	11.2	43.7		872,976	24.8	43.1	161,665	32.2		359,221	29.2		150,176	3.6	164,043	3.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6G
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by Metro Area

Comparison of Maternal Behavioral Indicators ^(2, 3)												
Metro Area	Medical Care					WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester			Pregpregnancy ⁽⁴⁾		During Preganancy ⁽⁵⁾		
	Number	% ^(Rank)	% ^(Rank)		Number	% ^(Rank)		Number	% ^(Rank)	Number	% ^(Rank)	
Dane Cty	1,889	83.9 (2)	1.7 (2)		2,058	25.5 (4)		1,841	18.5 (2)	1,537	84.5 (2)	
Mil. Cty	9,374	78.9 (4)	2.6 (4)		10,015	34.1 (2)		9,104	13.4 (3)	7,952	80.6 (3)	
Racine	1,292	81.1 (3)	1.3 (1)		1,430	39.2 (1)		1,254	10.8 (4)	1,128	78.2 (4)	
Rest of WI	18,730	85.5 (1)	2.3 (3)		20,272	34.0 (3)		17,611	20.1 (1)	15,531	85.5 (1)	
Wisconsin	31,285	83.3	2.3		33,775	33.7		29,810	17.6	26,148	83.6	
Nation (Prior Year)	579,816	78.5	5.6		944,971	31.2		38,592	15.7	206,967	74.1	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7G
2007 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by Metro Area

Comparison of Smoking Indicators (2, 3)										
Metro Area	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)		Number	% (Rank)	Number	% (Rank)	
Dane Cty	2,032	29.4 (2)	1,914	17.3 (2)	388	29.4 (2)	1,656	16.2 (1)	1,906	4.8 (1)
Mil. Cty	9,825	24.6 (1)	9,438	13.5 (1)	1,786	33.9 (1)	8,520	23.4 (4)	9,426	12.2 (4)
Racine	1,402	32.1 (3)	1,324	17.7 (3)	340	28.5 (3)	1,206	21.9 (2)	1,323	7.4 (2)
Rest of WI	19,835	41.9 (4)	18,899	26.4 (4)	6,368	26.2 (4)	16,574	23.1 (3)	18,879	7.6 (3)
Wisconsin	33,094	35.6	31,575	21.6	8,882	28.0	27,956	22.7	31,534	8.8
Nation (Prior Year)	847,411	27.2	692,984	16.8	156,923	35.0	514,582	22.5	443,951	24.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8G
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by Metro Area

Comparison of Infant Health Indicators (2, 3)										
Metro Area	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	1,834	7.7 (2)	8.7 (4)		1,844	8.8 (1)	1,599	2.9 (2)	1,873	77.7 (1)
Mil. Cty	8,896	9.6 (3)	6.0 (1)		9,072	11.6 (3)	7,577	4.0 (4)	9,050	63.8 (3)
Racine	1,293	9.7 (4)	7.0 (2)		1,294	12.4 (4)	1,096	3.1 (3)	1,300	59.2 (4)
Rest of WI	18,232	6.8 (1)	8.6 (3)		18,487	9.8 (2)	16,014	2.8 (1)	18,524	66.1 (2)
Wisconsin	30,255	7.8	7.8		30,697	10.4	26,286	3.2	30,747	65.8
Nation (Prior Year)	972,241	8.3	6.8		873,261	12.1	802,123	3.7	897,379	64.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3G
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Metro Area
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dane Cty	5,706	40.1	20.6	27.4	0.4	8.3	1.9	1.3
Mil. Cty	28,645	18.4	49.7	25.3	0.4	3.8	1.2	1.1
Racine	4,060	39.0	27.7	30.1	0.3	0.5	1.2	1.2
Rest of WI	56,951	73.4	3.0	13.6	3.1	4.5	1.2	1.2
Unknown Metro Areas	3	*	*	*	*	*	*	*
Wisconsin	95,365	53.4	19.1	18.6	2.0	4.3	1.2	1.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4G
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Metro Area
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dane Cty	5,706	0.3	5.8	11.9	62.1	18.5	1.3	0.1
Mil. Cty	28,645	0.7	9.0	14.0	59.1	16.0	1.2	0.0
Racine	4,060	0.6	8.2	14.7	60.9	14.5	1.1	0.0
Rest of WI	56,951	0.2	5.4	14.2	63.5	15.5	1.1	0.1
Unknown Metro Areas	3	*	*	*	*	*	*	*
Wisconsin	95,365	0.4	6.6	14.0	62.0	15.8	1.2	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5G
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by Metro Area
Combined 3 Years

Comparison of Maternal Health Indicators (2, 3)																						
Metro Area	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy								
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum													
	Number	% (Rank)	%		Number	% (Rank)	% (Rank)		Number	% (Rank)		% (Rank)	Number	% (Rank)		% (Rank)						
Dane Cty	5,217	11.2 (4)	42.0 (1)		3,376	29.1 (3)	40.8 (2)		1,072	30.3 (2)		3,521	21.8 (2)		3,209	4.9 (3)	3,218	5.6(1)				
Mil. Cty	27,666	8.7 (1)	49.6 (4)		20,650	26.9 (2)	42.3 (3)		4,529	40.9 (4)		22,612	34.2 (4)		16,584	4.4 (1)	16,737	7.1(3)				
Racine	3,756	10.3 (2)	48.7 (3)		3,316	37.3 (4)	38.7 (1)		633	31.1 (3)		3,174	22.0 (3)		1,684	6.5 (4)	1,714	9.3(4)				
Rest of WI	54,228	10.9 (3)	43.9 (2)		40,200	24.3 (1)	44.4 (4)		8,837	26.3 (1)		43,189	21.3 (1)		30,670	4.7 (2)	30,768	5.7(2)				
Unknown Metro Areas	3	*	*	*	*	*		3	*	*	*	*		2	*	*	*		0	*	*	*
Wisconsin	90,870	10.2	45.7		67,545	26.0	43.3		15,073	31.1		72,497	25.3		52,147	4.7		52,437	6.2			

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6G
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by Metro Area
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)														
Metro Area	Medical Care					WIC Enrollment			Multivitamin Consumption					
	1st Trimester		No Care			1st Trimester			Pregpregnancy ⁽⁴⁾			During Preganancy ⁽⁵⁾		
	Number	% (Rank)	% (Rank)			Number	% (Rank)		Number	% (Rank)		Number	% (Rank)	
Dane Cty	3,836	80.1 (2)	5.4 (2)			5,305	24.3 (4)		1,858	18.4 (2)		1,543	84.4 (2)	
Mil. Cty	19,500	75.9 (3)	5.5 (3)			26,676	32.6 (3)		9,212	13.5 (3)		7,970	80.6 (3)	
Racine	2,575	72.9 (4)	11.0 (4)			3,755	34.7 (1)		1,273	11.1 (4)		1,132	78.1 (4)	
Rest of WI	39,896	83.3 (1)	5.0 (1)			53,364	32.9 (2)		17,818	20.2 (1)		15,568	85.5 (1)	
Unknown Metro Areas	3	* *	* *			3	* *		0	* *		0	* *	
Wisconsin	65,810	80.5	5.4			89,103	32.4		30,161	17.7		26,213	83.6	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7G
2005 - 2007 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by Metro Area
Combined 3 Years

Comparison of Smoking Indicators (2, 3)															
Metro Area	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household									
	3 Months Prior to Pregnancy			Last 3 Months of Pregnancy		Prenatal			Postpartum						
	Number	% (Rank)		Number		% (Rank)		Number	% (Rank)		Number	% (Rank)			
Dane Cty	4,240	27.8	(2)	4,723	17.2	(2)	839	32.5	(1)	3,477	21.1	(1)	4,540	12.8	(1)
Mil. Cty	21,681	24.0	(1)	24,444	14.6	(1)	4,181	32.3	(2)	18,468	26.2	(2)	23,794	17.9	(3)
Racine	3,042	32.9	(3)	3,357	21.1	(3)	801	27.8	(3)	2,551	26.8	(3)	3,314	14.8	(2)
Rest of WI	43,364	43.0	(4)	49,385	27.5	(4)	15,491	26.8	(4)	36,179	30.6	(4)	47,087	18.3	(4)
Unknown Metro Areas	3	*	*	3	*	*	1	*	*	3	*	*	3	*	*
Wisconsin	72,330	36.0		81,912	22.8		21,313	28.2		60,678	28.5		78,738	17.7	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8G
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by Metro Area
Combined 3 Years

Comparison of Infant Health Indicators (2, 3)											
Metro Area	Birthweight					Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)								
	Number	% (Rank)	% (Rank)			Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	4,813	7.1 (2)	8.5 (3)			4,608	9.6 (1)	4,060	2.8 (2)	4,904	75.7 (1)
Mil. Cty	24,323	9.4 (4)	6.4 (1)			23,366	12.1 (4)	19,986	3.7 (4)	25,016	61.6 (3)
Racine	3,356	8.3 (3)	7.6 (2)			3,174	10.7 (3)	2,781	3.3 (3)	3,370	57.6 (4)
Rest of WI	49,218	7.0 (1)	8.6 (4)			47,392	10.2 (2)	41,676	2.7 (1)	50,022	64.8 (2)
Unknown Metro Areas	3	*	*	*	*	3	*	*	2	*	*
Wisconsin	81,713	7.8	7.9			78,543	10.7	68,505	3.1	83,315	64.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.